

# DRUG FACTS

Active ingredient(s)

Purpose

## Use(s)

- temporarily relieves these cold, sinus and flu symptoms:
- nasal and sinus congestion
- stuffy nose
- headache
- sore throat
- minor body aches and pains
- fever

## Warnings

### Do not use

### Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer and/or nasal decongestant
- right before or after heart surgery
- in a child who is taking a prescription monoamine oxidase inhibitor [MAOI] (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

### Ask a doctor before use if you have

### Ask a doctor before use if the child has

- problems or serious side effects from taking pain relievers, fever reducers, or nasal decongestants
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- not been drinking fluids
- lost a lot of fluid due to vomiting or diarrhea
- high blood pressure
- heart or kidney disease
- asthma
- thyroid disease
- diabetes
- taken a diuretic

### Ask a doctor or pharmacist before use if you are

### Ask a doctor or pharmacist before use if the child is

- taking any other drug containing an NSAID (prescription or nonprescription) and/or pseudoephedrine or any other nasal decongestant
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other drug

### When using this product

### When using this product

### do not exceed recommended dosage

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### Stop use and ask a doctor if

### Stop use and ask a doctor if

- the child feels faint, vomits blood, or has bloody or black stools. These are signs of stomach bleeding.
- stomach pain or upset gets worse or lasts
- the child does not get any relief within first day (24 hours) of treatment
- fever, pain, or nasal congestion gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- the child gets nervous, dizzy, or sleepless

### If pregnant or breast-feeding

### Keep out of reach of children

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- do not give longer than 10 days, unless directed by a doctor (see Warnings)
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- only use enclosed measuring cup
- if needed, repeat dose every **6 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

### Dosing Chart

Weight (lb)	Age (yrs)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24-47	2 to 5	1 tsp or 5 mL
48-95	6 to 11	2 tsp or 10 mL

## Other information

- store between 20-25°C (68-77°F)
- do not use if bottle wrap, or foil inner seal imprinted with "Safety Seal<sup>®</sup>" is broken or missing
- see bottom panel for lot number and expiration date

### Drug Facts

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)

- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

***Inactive ingredients***

acesulfame potassium, anhydrous citric acid, D&C yellow #10, FD&C red #40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum